

## HYALURONIDASE AFTER CARE INSTRUCTIONS

Hyaluronidase is an enzyme that breaks down HYALURONIC ACID dermal fillers. It can also break down some of the naturally occurring hyaluronic acid present in the tissues of the body.

Results can be unpredictable. Volume can be lost dramatically when it is administered.

Results can be seen immediately although the final results may take up to two weeks.

The procedure may need to be repeated to ensure satisfactory results.

Administration of hyaluronidase can cause ANAPHYLAXIS (severe allergic reaction). This may be life threatening & requires IMMEDIATE MEDICAL ATTENTION. Symptoms include:

- Shortness of breath
- Wheezing
- Coughing
- Difficulty in swallowing
- Swollen tongue/eyelids/lips
- Hoarse voice
- Stomach pain
- Nausea or diarrhoea

IF YOU EXPERIENCE ANY OF THESE SYMPTOMS CONTACT EMERGENCY MEDICAL SERVICES IMMEDIATELY

Following your procedure other mild/moderate injection-related reactions may occur at the site of the injections. These include:

- Redness
- Swelling
- Pain
- Itching
- Bruising
- Tenderness
- 

These generally resolve spontaneously after a few days. If you are concerned about any of the above, contact your practitioner straight away.

As multiple treatments are typically required you will be asked to return for a review within 2-3 weeks. It is important that you attend. Let your practitioner know if you are unable to keep your appointment.

NEXT APPOINTMENT: .....



.....